# Psychological Science

## It's Not All About Me : Motivating Hand Hygiene Among Health Care Professionals by Focusing on Patients

Adam M. Grant and David A. Hofmann Psychological Science 2011 22: 1494 originally published online 10 November 2011 DOI: 10.1177/0956797611419172

> The online version of this article can be found at: http://pss.sagepub.com/content/22/12/1494

> > Published by: SAGE http://www.sagepublications.com

> > > On behalf of:



Association for Psychological Science

Additional services and information for *Psychological Science* can be found at:

Email Alerts: http://pss.sagepub.com/cgi/alerts

Subscriptions: http://pss.sagepub.com/subscriptions

Reprints: http://www.sagepub.com/journalsReprints.nav

Permissions: http://www.sagepub.com/journalsPermissions.nav

>> Version of Record - Dec 8, 2011 Proof - Nov 10, 2011 What is This?

### It's Not All About Me: Motivating Hand Hygiene Among Health Care Professionals by Focusing on Patients

Psychological Science 22(12) 1494–1499 © The Author(s) 2011 Reprints and permission: sagepub.com/journalsPermissions.nav DOI: 10.1177/0956797611419172 http://pss.sagepub.com

PSYCHOLOGICAL SCIENCE



#### Adam M. Grant<sup>1</sup> and David A. Hofmann<sup>2</sup>

<sup>1</sup>Management Department, The Wharton School, University of Pennsylvania, and <sup>2</sup>Kenan-Flagler Business School, University of North Carolina at Chapel Hill

#### Abstract

Diseases often spread in hospitals because health care professionals fail to wash their hands. Research suggests that to increase health and safety behaviors, it is important to highlight the personal consequences for the actor. However, because people (and health care professionals in particular) tend to be overconfident about personal immunity, the most effective messages about hand hygiene may be those that highlight its consequences for other people. In two field experiments in a hospital, we compared the effectiveness of signs about hand hygiene prevents patients from catching diseases") or patient safety ("Hand hygiene prevents patients from catching diseases"). We assessed hand hygiene by measuring the amount of soap and hand-sanitizing gel used from dispensers (Experiment 1) and conducting covert, independent observations of health care professionals' hand-hygiene behavior: (Experiment 2). Results showed that changing a single word in messages motivated meaningful changes in behavior: The hand hygiene of health care professionals increased significantly when they were reminded of the implications for patients but not when they were reminded of the implications for themselves.

#### **Keywords**

social influences, motivation, health, cooperation

Received 5/27/11; Revision accepted 7/12/11

In 1847, Ignaz Semmelweis required health care professionals at the Vienna General Hospital to wash their hands, and death rates due to childbed fever decreased from 18.3% to 1.3%. Since then, extensive research has demonstrated that hand hygiene plays a critical role in preventing the spread of infections and diseases (Backman, Zoutman, & Marck, 2008). Nevertheless, it is common for health care professionals to wash their hands less than half as often as recommended, and many interventions for improving hand hygiene among health care professionals have proven ineffective (Gawande, 2004; Whitby et al., 2007). How can psychological science guide the development of messages to address this pressing problem?

Messages about health and safety are thought to be effective when they highlight personal risks for the actor. Researchers have speculated that health care professionals "are probably driven to wash their hands by their need to protect themselves more than [by their need to protect] their patients" (Korniewicz & El-Masri, 2010, p. 88). According to this line of logic, messages aimed at health care professionals should emphasize how hand hygiene protects them personally. Such messages are believed to activate basic motivations related to survival and self-protection (Rothman & Salovey, 1997). As Williams and Noyes (2007) summarized, safety behavior "is dependent on individuals believing that the risk is likely to affect *them*, that it will have serious consequences for *them*" (p. 21, emphasis in original).

However, research on overconfidence has shown that individuals consistently overestimate their immunity (Dunning, Heath, & Suls, 2004). This illusion of invulnerability is common among health care professionals. As two physicians explained, "I'm a doctor, I'm protected," and "We doctors wear magic white coats. We destroy disease. . . . How could it ever attack us?" (Klitzman, 2006, p. 547).

Overconfidence is likely to be fueled by both motivational and cognitive processes. First, to maintain a sense of security while working in hazardous environments, health care professionals may need to convince themselves that they are protected. According to research on motivated reasoning (Kunda, 1990) and confirmation biases (Nickerson, 1998), health care professionals may search for information that seems to verify their personal safety and may discount information that

**Corresponding Author:** Adam M. Grant, University of Pennsylvania, 3620 Locust Walk, Suite 2000 SH/DH, Philadelphia, PA 19104-6370 E-mail: grantad@wharton.upenn.edu suggests they are at risk. Indeed, research has shown that people tend to respond defensively to information that poses a threat to their personal health or safety: People are likely to scrutinize such messages for flaws rather than accept the information they contain (Liberman & Chaiken, 1992).

Second, although health care professionals are frequently exposed to diseases, they contract relatively few. When they do get sick, it is not clear that poor hand hygiene is the culprit. Thus, it may be easy for health care professionals to recall instances in which they failed to wash their hands without getting sick, but difficult for them to recall episodes in which failing to wash their hands made them ill. According to research on the availability heuristic (Tversky & Kahneman, 1974), because the ease with which an event comes to mind serves as a cue for its likelihood (Schwarz et al., 1991), health care professionals should perceive that failing to wash their hands poses little personal risk. Consequently, messages emphasizing the personal consequences of hand hygiene for health care professionals may fall on deaf ears.

Research on persuasion reveals that for a message to resonate with an audience, it must be relevant to that audience's perspective (Cialdini, 2003; Clary & Snyder, 1999; Rothman & Salovey, 1997). We hypothesized that health care professionals would be more motivated to wash their hands by messages highlighting patient consequences than by messages highlighting personal consequences. Whereas people tend to overestimate their own invulnerability, for both motivational and cognitive reasons, they are less susceptible to this bias when estimating the vulnerability of other people (Dunning et al., 2004). Moreover, the fact that patients are by definition a vulnerable population may make their risks salient to health care professionals, who are trained to err in favor of caution when treating patients (Gigerenzer, Gaissmaier, Kurz-Milcke, Schwartz, & Woloshin, 2008).

Thus, messages aimed at health care professionals should be most effective when they emphasize how hand-hygiene practices can protect patients' health rather than personal health. We tested this hypothesis in two field experiments in a hospital by subtly manipulating the content of signs about hand hygiene and testing their influence with unobtrusive measures (Webb, Campbell, Schwartz, & Sechrest, 1966). To measure the signs' effects, we used two strong, complementary assessment techniques recommended by The Joint Commission (2009) as part of the Consensus Measurement in Hand Hygiene project: objective measurements of the use of hand-hygiene products and independent observations of adherence to safe hand-hygiene practices.

#### Experiment I Method

We compared the effects of signs emphasizing personal consequences, patient consequences, or neither on the hand-hygiene behaviors of health care professionals in a U.S. hospital. Our focus on signs was based on evidence that small variations in the content of messages can produce powerful changes in mindsets and behaviors (Cialdini, 2003; Crum & Langer, 2007). We assessed hand hygiene by measuring the percentage of soap and hand-sanitizing gel used from dispensers in hospital units; this technique was both objective and unobtrusive (The Joint Commission, 2009). Our sample comprised 66 dispensers available for physicians and nurses in the hospital, and we measured the amount of soap and gel used during 2-week periods before and after we introduced our signs. To measure baseline product use, we had an environmental-services team fill the bag in each dispenser with soap or gel and weigh each bag 2 weeks later. This team, which was blind to our hypotheses, then refilled the dispensers before we began the experiment. To minimize demand characteristics and cross-contamination, we did not inform employees at the hospital that research was underway.

We randomly assigned one of three signs to each dispenser. The personal-consequences sign read, "Hand hygiene prevents you from catching diseases." The patient-consequences sign read, "Hand hygiene prevents patients from catching diseases." The control sign, which was developed by hospital managers, read, "Gel in, wash out." Except for these subtle differences in wording, the signs were identical. One sign was posted above each dispenser by a safety professional. After 2 weeks, the environmental-services team weighed each bag again. Because the bags were of different sizes, we report our dependent measure as the percentage by weight of soap or gel used.

#### **Results and discussion**

Table 1 reports the mean percentage of soap or gel used from the dispensers in each condition, before and after the signs were introduced. A repeated measures analysis of variance on the amount of hand-hygiene product used showed a significant interaction between time (pretest, posttest) and condition

 Table 1. Mean Percentage (by Weight) of Hand-Hygiene Products Used as a Function of Condition in Experiment 1

Condition	Pretest product usage	Posttest product usage		
Control $(n = 21)$	38.24% (24.90)	40.13% (24.43)		
Personal consequences $(n = 23)$	35.49% (28.18)	33.98% (19.65)		
Patient consequences $(n = 22)$	37.25% (36.46)	54.18% (18.33)		

Note: Standard deviations are in parentheses.

(personal consequences, patient consequences, control), F(2, 63) = 3.30, p = .04,  $\eta^2 = .09$ ,  $p_{rep} = .89$ . Paired-samples *t* tests showed a significant increase from pretest to posttest in the amount of hand-hygiene product used from dispensers with the patient-consequences sign (37.25% to 54.18%), t(21) = 2.72, p = .01, d = 0.59,  $p_{rep} = .96$ , but not from dispensers with the personal-consequences sign (35.49% to 33.98%), t(22) = -0.27, p = .79, or the control sign (38.24% to 40.13%), t(20) = 0.64, p = .53. Tukey's multiple comparison test showed no significant differences between conditions in the pretest use of hand-hygiene product, but in the posttest, the amount of hand-hygiene product used from dispensers with the patient-consequences sign was significantly greater than the amount used from dispensers with the personal-consequences sign (p < .01, d = 1.06,  $p_{rep} = .97$ ) or the control sign (p < .05, d = 0.65,  $p_{rep} = .89$ ).

Although these results are encouraging, they are subject to two key limitations. First, health care professionals may have been influenced by a sign near one dispenser but used handhygiene products from a different dispenser. Second, it is possible that the effects were influenced either by a small number of health care professionals who used large quantities of handhygiene products or by patients who gained access to the dispensers. To address these potential confounds, in Experiment 2, we assigned the personal-consequences and patientconsequences signs to separate hospital units and asked experts to directly observe health care professionals' handhygiene behaviors.

#### Experiment 2 Method

Nine months after Experiment 1, we conducted a second experiment in different units of the same hospital, using the same personal- and patient-consequences signs. We enlisted three experts—a physician in charge of patient safety, an infectious-disease specialist, and a lead nurse manager to organize hospital units into matched pairs on the basis of similar types of patients, health conditions, and professional specialties. The three experts achieved consensus on four matched pairs of units: pediatric and neonatal intensive care units (ICUs), cardiac and neurological critical care units (CCUs), cardiology and chest-pain units, and generalobservation and medical-teaching units. We assigned the personal-consequences sign to four of the units (pediatric ICU, cardiac CCU, cardiology, and general observation) and the patient-consequences sign to the other four units (neonatal ICU, neurological CCU, chest pain, and medical teaching).

We measured hand hygiene using observations of the behaviors of health care professionals. For each unit, the patient-safety team identified one expert observer with professional training and certifications in nursing. The observers were blind to our hypotheses and conducted their observations covertly, thereby minimizing demand characteristics and reducing the likelihood that the observed health care professionals would be aware that research was underway and that their behavior was being tracked. Following the guidelines recommended by the Consensus Measurement in Hand Hygiene project team (The Joint Commission, 2009), we asked the observers to count health care professionals' opportunities for hand hygiene and to indicate whether the health care professional in each instance adhered to safe handhygiene practices. We defined an opportunity for hand hygiene as occurring before or after contact with a patient. To ensure that only health care professionals' behavior was included in the data, the observers recorded the date of each hand-hygiene opportunity and the type of practitioner involved. Observers identified three types of practitioners: nurses (59% of observations), physicians (17% of observations), and ancillary staff (technicians, nutritionists, social workers, pharmacists, and transporters; 24% of observations).

We collected pretest data over a 2-week period, during which the observers identified 322 hand-hygiene opportunities (the practitioners adhered to hand-hygiene guidelines in 259 of these instances). After the pretest, medical-safety professionals posted the signs in their assigned units. The ratios of signs to patient beds were equivalent in the two conditions: .80 for units assigned to the personal-consequences condition (57 signs, 71 beds) and .79 for units assigned to the patient-consequences condition (69 signs, 87 beds). The observers identified 245 hand-hygiene opportunities during the 2-week posttest period (the practitioners adhered to hand-hygiene guidelines in 212 of these instances). We tested whether hand-hygiene adherence increased on units with the patient-consequences sign but not on units with the personal-consequences sign.

#### **Results and discussion**

Table 2 reports the percentage of instances in which health care practitioners adhered to safe hand-hygiene practices in each condition, before and after the signs were introduced. A contingency-table analysis showed that hand-hygiene adherence increased

**Table 2.** Adherence to Safe Hand-Hygiene Practices as a Function of Condition inExperiment 2

Condition	Pretest hand-hygiene adherence	Posttest hand-hygiene adherence		
Personal consequences	80.00% (96/120)	79.71% (55/69)		
Patient consequences	80.69% (163/202)	89.20% (157/176)		

significantly on units with the patient-consequences sign (80.69% to 89.20%),  $\chi^2(1, N = 378) = 5.25$ , p = .02, d = 0.33,  $p_{\rm rep} = .93$ . In contrast, hand-hygiene adherence did not change significantly on units with the personal-consequences sign (80.00% to 79.71%),  $\chi^2(1, N = 189) = 0.04$ , p = .85. During the pretest period, the units assigned to the personal- and patient-consequences conditions did not differ significantly in hand-hygiene adherence,  $\chi^2(1, N = 322) = 0.02$ , p = .88. However, during the posttest period, hand-hygiene adherence was significantly greater on units with the patient-consequences sign than on units with the personal-consequences sign,  $\chi^2(1, N = 245) = 3.83$ , p = .05, d = 0.36,  $p_{\rm rep} = .88$ .

Although the units were matched and then randomly assigned to condition, we reanalyzed the data to control for unit; the same pattern of results emerged. To examine whether the observed effects of the patient-consequences sign were due to unique characteristics of the high-risk units or of the health care professionals who worked in such units, we compared the higher-risk units (ICUs and CCUs) with the lower-risk units. Binary logistic regression analyses showed no significant differences between unit types in the effects of the patientconsequences sign (b = 0.31, SE = 0.62, Wald z = 0.25, p = .62) or of the personal-consequences sign (b = 0.19, SE = 0.88, Wald z = 0.05, p = .83; this result suggests that the effects were robust across units. We also investigated whether the effects varied by practitioner type (see Table 3). The patientconsequences sign significantly increased hand hygiene for physicians, marginally increased it for nurses, and did not affect the hand-hygiene behavior of ancillary staff. The personalconsequences sign had no significant effects.

#### **General Discussion**

Together, these findings suggest that messages about patient consequences, rather than personal consequences, can encourage hand hygiene among health care professionals. Our results have important theoretical and practical implications for the design of persuasive communications about health and safety. In theoretical terms, whereas research has typically focused on the effects of highlighting the personal consequences of health- and safety-related behaviors (Williams & Noyes, 2007), our studies demonstrate the value of highlighting the consequences of such behaviors for other people. Psychologists have long recognized that seemingly innocuous situational forces, such as time pressure, can impede prosocial behaviors even among good Samaritans with the best of intentions (Darley & Batson, 1973). It is tempting to conclude that capturing the attention of busy health care professionals to encourage hand hygiene depends on appealing to their immediate self-interest (Korniewicz & El-Masri, 2010). However, our research reveals that reminders of prosocial consequences may have a greater influence on the hand-hygiene behavior of health care professionals than reminders of personal consequences do (see also Grant, 2008).

In practical terms, the significant effects of our subtle experimental manipulation on a difficult-to-change dependent variable have substantial real-world implications (Prentice & Miller, 1992). Over 2-week periods, the patient-consequences signs produced an increase of more than 45% in the amount of hand-hygiene product used per dispenser (Experiment 1) and an increase of more than 10% in hand-hygiene behavior among health care professionals before and after contact with patients (Experiment 2). These results are particularly meaningful given that the few hand-hygiene interventions known to be successful tend to rely on expensive technologies and large-scale cultural changes (Pittet et al., 2000; Whitby et al., 2007).

A key limitation of both studies is that they lasted for only 2 weeks. Because the effects of hand-hygiene interventions are often short-lived (Pittet et al., 2000), an examination of their sustainability is of critical importance. If they fade because of habituation and desensitization, researchers should explore strategies for maintaining novelty, such as rotating messages or incorporating photos of patients into messages. Nevertheless, 2 weeks of increased adherence to safe hand-hygiene practices can have considerable effects. Applying the findings of Pittet et al. (2000) and Rosenthal, Guzman, and Safdar (2005), we estimated the number of infections prevented in the patient-consequences condition in each study to

Table 3. Results of Analyses of Adherence to Safe Hand-Hygiene Practices in Experiment 2

<b>T C</b>	Hand-hygiene adherence					
Type of practitioner and condition	Pretest	Posttest	$\chi^2(1)$	Þ	d	₽ <sub>rep</sub>
Nurses						
Personal consequences	81.43% (57/70)	83.87% (26/31)	0.09	.77	_	
Patient consequences	79.53% (101/127)	88.39% (99/112)	3.43	.06	0.33	.86
Physicians	, , , , , , , , , , , , , , , , , , ,	. ,				
Personal consequences	72.73% (16/22)	87.50% (14/16)	1.22	.27	_	
Patient consequences	72.41% (21/29)	92.86% (26/28)	4.12	.04	0.76	.89
Ancillary staff						
Personal consequences	85.19% (23/27)	68.18% (15/22)	-2.01	.16	_	_
Patient consequences	89.13% (41/46)	88.89% (32/36)	-0.00	.97	—	—

be between two and nine; these infections would have cost the hospital between \$9,000 and \$30,000 per study (see Chen, Chou, & Chou, 2005). This prevention of infections is a substantial return on investment, given the minimal costs of printing and posting signs. If the increased hand-hygiene adherence were sustained for a year across the hospital, the potential benefits could include the prevention of more than 100 infections and a savings of more than \$300,000.

Future research should test perceived vulnerability and other mediating mechanisms. For example, the patient-consequences sign highlighted the implications for a group, whereas the personal-consequences sign highlighted the implications for an individual. Did the responsiveness to messages about patient consequences reflect utilitarian reasoning, whereby health care professionals aimed to promote the greatest good for the greatest number of people? Given that individuals are more likely to help a single person than to help multiple people (Kogut & Ritov, 2007; Slovic, 2007), the opposite seems likely. Signs might catalyze greater empathy by mentioning "a patient" or "the patient in this room" instead of "patients" (Goldstein, Cialdini, & Griskevicius, 2008). However, the effects of this change in wording might be mitigated by health care professionals' perception of patients as part of a unitary, cohesive group (Burson, Smith, & Faro, 2010). To investigate these issues, future studies should systematically vary group size. Nevertheless, if group size is a contributing factor, it may be an explanatory mechanism rather than a confound: Handhygiene behaviors can protect only one self, but many other people. This fact accentuates the value of examining whether patients are perceived as being part of a uniquely vulnerable and valued population. Would similar effects emerge if signs highlighted other groups affected by hand hygiene, such as nurses, physicians, or health care professionals in general?

We also recommend combining quantity-based measures of the use of hand-hygiene products and frequency-based observational measures with quality-based measures, such as microbiological tests and assessments of infection rates. Finally, future research should investigate the implications of our findings for other health, safety, and prosocial behaviors in different populations. Are people more likely to improve their exercise and eating habits, to quit smoking, to purchase life insurance, to wear seat belts and helmets, to protect the environment, or to take prescription medication when they are reminded of the consequences of these behaviors for their families rather than for themselves?

In conclusion, our findings suggest that health and safety messages should focus not on the self, but rather on the target group that is perceived as most vulnerable. As Levitt and Dubner (2009) suggested, "When a doctor fails to wash his own hands, his own life isn't the one that is primarily endangered. [The life endangered is that of] the next patient he treats" (p. 207). Merely emphasizing the consequences for patients motivates health care professionals to take more everyday healthprotective action. From the perspective of a health care professional, safety behavior is not necessarily "all about me."

#### Acknowledgments

The authors thank Jessica Dixon, Janine Jones, Meera Kelley, and Betty Woodard for assistance with data collection and Associate Editor Julie Fitness, Noah Eisenkraft, Francesca Gino, and two anonymous reviewers for feedback.

#### **Declaration of Conflicting Interests**

The authors declared that they had no conflicts of interest with respect to their authorship or the publication of this article.

#### References

- Backman, C., Zoutman, D. E., & Marck, P. B. (2008). An integrative review of the current evidence on the relationship between hand hygiene interventions and the incidence of health care-associated infections. *American Journal of Infection Control*, 36, 333–348.
- Burson, K., Smith, R., & Faro, D. (2010, November). *The influence of entitativity on charitable giving*. Paper presented at the annual meeting of the Society for Judgment and Decision Making, St. Louis, MO.
- Chen, Y. Y., Chou, Y. C., & Chou, P. (2005). Impact of nosocomial infection on cost of illness and length of stay in intensive care units. *Infection Control & Hospital Epidemiology*, 26, 281–287.
- Cialdini, R. B. (2003). Crafting normative messages to protect the environment. *Current Directions in Psychological Science*, 12, 105–109.
- Clary, E. G., & Snyder, M. (1999). The motivations to volunteer: Theoretical and practical considerations. *Current Directions in Psychological Science*, 8, 156–159.
- Crum, A. J., & Langer, E. J. (2007). Mind-set matters: Exercise and the placebo effect. *Psychological Science*, 18, 165–171.
- Darley, J. M., & Batson, C. D. (1973). From Jerusalem to Jericho: A study of situational and dispositional variables in helping behavior. *Journal of Personality and Social Psychology*, 27, 100–108.
- Dunning, D., Heath, C., & Suls, J. M. (2004). Flawed self-assessment: Implications for health, education, and the workplace. *Psychological Science in the Public Interest*, 5, 69–106.
- Gawande, A. (2004). Notes of a surgeon: On washing hands. *New England Journal of Medicine*, *350*, 1283–1286.
- Gigerenzer, G., Gaissmaier, W., Kurz-Milcke, E., Schwartz, L. M., & Woloshin, S. (2008). Helping doctors and patients make sense of health statistics. *Psychological Science*, *8*, 53–96.
- Goldstein, N. J., Cialdini, R. B., & Griskevicius, V. (2008). A room with a viewpoint: Using social norms to motivate environmental conservation in hotels. *Journal of Consumer Research*, 35, 472–482.
- Grant, A. M. (2008). The significance of task significance: Job performance effects, relational mechanisms, and boundary conditions. *Journal of Applied Psychology*, 93, 108–124.
- The Joint Commission. (2009). Measuring hand hygiene adherence: Overcoming the challenges. Retrieved from http://www .jointcommission.org/Measuring\_Hand\_Hygiene\_Adherence\_ Overcoming\_the\_Challenges\_/
- Klitzman, R. (2006). Post-residency disease and the medical self: Identity, work, and health care among doctors who become patients. *Perspectives in Biology and Medicine*, 49, 542–552.

- Kogut, T., & Ritov, I. (2007). "One of us": Outstanding willingness to help save a single identified compatriot. Organizational Behavior and Human Decision Processes, 104, 150–157.
- Korniewicz, D. M., & El-Masri, M. (2010). Exploring the factors associated with hand hygiene compliance of nurses during routine clinical practice. *Applied Nursing Research*, 23, 86–90.
- Kunda, Z. (1990). The case for motivated reasoning. *Psychological Bulletin*, 108, 480–498.
- Levitt, S. D., & Dubner, S. J. (2009). Superfreakonomics: Global cooling, patriotic prostitutes, and why suicide bombers should buy life insurance. New York, NY: HarperCollins.
- Liberman, A., & Chaiken, S. (1992). Defensive processing of personally relevant health messages. *Personality and Social Psychology Bulletin*, 18, 669–679.
- Nickerson, R. S. (1998). Confirmation bias: A ubiquitous phenomenon in many guises. *Review of General Psychology*, 2, 175–220.
- Pittet, D., Hugonnet, S., Harbarth, S., Mourouga, P., Sauvan, V., Touveneau, S., & Perneger, T. (2000). Effectiveness of a hospitalwide programme to improve compliance with hand hygiene. *The Lancet*, 356, 307–312.
- Prentice, D. A., & Miller, D. T. (1992). When small effects are impressive. *Psychological Bulletin*, 112, 160–164.
- Rosenthal, V. D., Guzman, S., & Safdar, N. (2005). Reduction in nosocomial infection with improved hand hygiene in intensive care

units of a tertiary care hospital in Argentina. *American Journal of Infection Control*, 33, 392–397.

- Rothman, A. J., & Salovey, P. (1997). Shaping perceptions to motivate healthy behavior: The role of message framing. *Psychologi*cal Bulletin, 121, 3–19.
- Schwarz, N., Bless, H., Strack, F., Klumpp, G., Rittenauer-Schatka, H., & Simons, A. (1991). Ease of retrieval as information: Another look at the availability heuristic. *Journal of Personality* and Social Psychology, 61, 195–202.
- Slovic, P. (2007). "If I look at the mass I will never act": Psychic numbing and genocide. Judgment and Decision Making, 2, 79–95.
- Tversky, A., & Kahneman, D. (1974). Judgment under uncertainty: Heuristics and biases. *Science*, 185, 1124–1131.
- Webb, E. J., Campbell, D. T., Schwartz, R. D., & Sechrest, L. (1966). Unobtrusive measures: Nonreactive research in the social sciences. Chicago, IL: Rand McNally.
- Whitby, M., Pessoa-Silva, C. L., McLaws, M.-L., Allegranzi, B., Sax, H., Larson, E., . . . Pittet, D. (2007). Behavioural considerations for hand hygiene practices: The basic building blocks. *Journal of Hospital Infection*, 65, 1–8.
- Williams, D. J., & Noyes, J. M. (2007). How does our perception of risk influence decision-making? Implications for the design of risk information. *Theoretical Issues in Ergonomics Science*, 8, 1–35.